

SERI/TR-642-761

February 1981

# A Simplified Clear Sky Model for Direct and Diffuse Insolation on Horizontal Surfaces

Richard E. Bird  
Roland L. Hulstrom



# SERI

**Solar Energy Research Institute**

A Division of Midwest Research Institute

1617 Cole Boulevard  
Golden, Colorado 80401

Operated for the  
**U.S. Department of Energy**  
under Contract No. EG-77-C-01-4042

Printed in the United States of America

Available from:

National Technical Information Service

U.S. Department of Commerce

5285 Port Royal Road

Springfield, VA 22161

Price:

Microfiche \$3.00

Printed Copy \$ 5.25

## NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

SERI/TR-642-761

UC CATEGORIES: UC-59,61,62,63

A SIMPLIFIED CLEAR SKY MODEL  
FOR DIRECT AND DIFFUSE INSOLATION  
ON HORIZONTAL SURFACES

RICHARD E. BIRD  
ROLAND L. HULSTROM

FEBRUARY 1981

PREPARED UNDER TASK NO. 1093.00

**Solar Energy Research Institute**

A Division of Midwest Research Institute

1617 Cole Boulevard  
Golden, Colorado 80401

Prepared for the  
U.S. Department of Energy  
Contract No. EG-77-C-01-4042

**PREFACE**

The work documented here was performed by the SERI Renewable Resource Assessment Branch for the U.S. Department of Energy under Task No. 1093.00. The report compares several simple global horizontal insolation models with several rigorous radiative transfer models and describes an improved, simple, global insolation model. We would like to thank J. V. Dave of IBM for providing data sets from his Spherical Harmonics code.

*Richard E Bird*

---

Richard E. Bird  
Senior Scientist

Approved for

SOLAR ENERGY RESEARCH INSTITUTE

*Roland L. Hulstrom*

---

Roland L. Hulstrom, Chief  
Renewable Resource Assessment Branch

*Barry L Butler*

---

Barry Butler, Manager  
General Research Division

**TABLE OF CONTENTS**

	<u>Page</u>
1.0 Introduction.....	1
2.0 Description of Models.....	3
2.1 Atwater and Ball Model.....	3
2.2 Davies and Hay Model.....	4
2.3 Watt Model.....	5
2.4 Hoyt Model.....	6
2.5 Lacis and Hansen Model.....	7
2.6 Bird Model.....	7
2.7 Rigorous Codes.....	10
3.0 Model Comparisons.....	13
3.1 Transmittance Comparisons.....	13
3.2 Irradiance Comparisons.....	15
4.0 Summary and Conclusions.....	27
5.0 References.....	29
Appendix A: Tabulated Model Data.....	31

**LIST OF FIGURES**

	<u>Page</u>
3-1 Transmittance versus Secant of Solar Zenith Angle for Midlatitude Summer Model.....	14
3-2 Aerosol Transmittance--USS Atmosphere (23-km-Visibility Aerosol)....	16
3-3 Rayleigh Transmittance.....	16
3-4 Water Vapor Transmittance--USS Atmosphere (1.42-cm H <sub>2</sub> O in Vertical Column).....	17
3-5 Ozone Transmittance--USS Atmosphere.....	17
3-6 Total Insolation--USS Atmosphere.....	18
3-7 Direct Horizontal Insolation--USS Atmosphere.....	18
3-8 Diffuse Sky Insolation--USS Atmosphere.....	19
3-9 Diffuse Ground/Sky Insolation--USS Atmosphere.....	19
3-10 Total Insolation--MLS Atmosphere.....	21
3-11 Direct Horizontal Insolation--MLS Atmosphere.....	21
3-12 Diffuse Sky Insolation--MLS Atmosphere.....	22
3-13 Diffuse Ground/Sky Insolation--MLS Atmosphere.....	22
3-14 Total Insolation--Dave Model 3.....	23
3-15 Direct Horizontal Insolation--Dave Model 3.....	23
3-16 Diffuse Sky Insolation--Dave Model 3.....	24
3-17 Diffuse Ground/Sky Insolation--Dave Model 3.....	24
3-18 Direct Normal Insolation--Dave Model 3.....	26
3-19 Direct Normal Insolation--USS Atmosphere.....	26

## LIST OF TABLES

	<u>Page</u>
2-1 Equations for Total Downward Irradiance for Atwater and Ball Model.....	3
2-2 Equations for Total Downward Irradiance for Davies and Hay Model....	4
2-3 Equations for Total Downward Irradiance for Watt Model.....	5
2-4 Equations for Total Downward Irradiance for Hoyt Model.....	6
2-5 Equations for Total Downward Irradiance for Lacis and Hansen Model.....	7
2-6 Equations for Total Downward Irradiance for the Bird Model.....	8
2-7 Values of the Asymmetry Factor for the Rural and the Dave Haze L Aerosol.....	9
2-8 Input Parameters Required for Simple Models.....	10
3-1 Amounts of H <sub>2</sub> O and O <sub>3</sub> in a Vertical Column for the MLS and USS Atmospheres.....	13
A-1 Tabulated Data from Several Models for the USS Atmosphere.....	33
A-2 Tabulated Data from Several Models for the MLS Atmosphere.....	35
A-3 Tabulated Data from Several Models for the Dave Model 3 Atmosphere.....	37

## NOMENCLATURE

$a_i$	Absorptance parameters defined by Hoyt
$a_{CO}$	Carbon dioxide absorptance
$a_{OX}$	Oxygen absorptance
$a_W$	Water vapor absorptance
$B_a$	Ratio of the forward-scattered irradiance to the total scattered irradiance due to aerosols
$F_{zj}$	A parameter used by Watt in air mass calculations
$I_{as}$	Solar irradiance on a horizontal surface from atmospheric scattering ( $W/m^2$ )
$I_d$	Direct solar irradiance on a horizontal surface ( $W/m^2$ )
$I_G$	Solar irradiance on a horizontal surface from multiple reflections between the ground and sky ( $W/m^2$ )
$I_s$	Solar irradiance on a horizontal surface from scattered light ( $I_{as} + I_G$ )
$I_T$	Total (global) solar irradiance on a horizontal surface ( $W/m^2$ )
$I_o$	Extraterrestrial solar irradiance ( $1353 W/m^2$ )
$K$	Aerosol transmittance for Davies and Hay
$K_1$	Constant used in Bird model associated with aerosol absorptance
$M$	Air mass
$M'$	Pressure-corrected air mass
$M_i$	Watt's path length modifier (similar to air mass)
$P$	Surface pressure (millibars)
$r_g$	Ground albedo
$r_s$	Sky, or atmospheric, albedo
$T$	Surface temperature (K)
$T_A$	Transmittance of aerosol absorptance and scattering
$T_{AA}$	Transmittance of aerosol absorptance

$T_{AS}$	Transmittance of aerosol scattering
$T_{as}$	Transmittance of dry air absorptance and scattering for Watt
$T_L$	Transmittance of lower layer aerosol for Watt
$T_M$	Global transmittance of all molecular effects except water vapor for Atwater
$T_{Md}$	Direct transmittance of all molecular effects except water vapor for Atwater
$T_O$	Transmittance of ozone absorptance
$T_R$	Transmittance of Rayleigh scattering
$T_U$	Transmittance of upper layer aerosol for Watt
$T_{UM}$	Transmittance of absorptance of uniformly mixed gases (carbon dioxide and oxygen)
$T_w$	Transmittance of water vapor absorptance ( $1 - a_w$ )
$T_{ws}$	Transmittance of water vapor scattering
$U_O$	Amount of ozone in a vertical column from surface (cm)
$U_w$	Amount of precipitable water in a vertical column from surface (cm)
$X_O$	Total amount of ozone in a slanted path (cm)
$X_w$	Total amount of precipitable water in a slanted path (cm)
$Z$	Angle between a line to the sun and the local zenith (zenith angle in degrees)
$\alpha_{cs}$	An attenuation multiplier used by Watt
$\tau_A$	Broadband aerosol optical depth from surface in a vertical path (broadband turbidity)
$\tau_{0.5}$	Aerosol optical depth from surface in a vertical path at 0.5- $\mu$ m wavelength
$\tau_{0.38}$	Aerosol optical depth from surface in a vertical path at 0.38- $\mu$ m wavelength
$\tau_L$	Lower layer aerosol optical depth in a vertical path for Watt
$\tau_u$	Upper layer aerosol optical depth in a vertical path for Watt
$W_O$	Single scattering albedo--the fraction of the light lost from an incident pencil of radiation that is due to scattering

## SECTION 1.0

### INTRODUCTION

To properly design a solar energy system for a location lacking an insolation data base, insolation models are required. This need for insolation models has been recognized for many years. One early and widely used insolation model was published in 1940 by Moon [1]. This model is still used today in its original or modified forms [2,3].

Insolation models have proliferated to the point where it is difficult for a solar user to decide which model to adopt. The purpose of this paper is to provide a detailed comparison of several simple, broadband insolation models that are currently in use and, based on this comparison and comparisons with more rigorous radiative transfer models, to formulate a simple clear sky model for direct and diffuse insolation. This type of comparison should be helpful in evaluating the relative accuracy of various models and give direction for formulating a model that uses the best of each existing model. The criteria used for evaluating and formulating models have been simplicity, accuracy, and the ability to use readily available meteorological data.

It should be noted that the use of the word "rigorous" does not necessarily mean that the "rigorous" results truly represent reality. Even though these "rigorous" codes are very detailed in the methods used to solve the radiative transfer problem, the representativeness of the results depends upon how well the atmospheric model, the measured atmospheric parameters, the mathematical methods, and other assumptions made in the codes relate to a real situation. However, in the absence of well-documented data, this approach of using "rigorous" codes as a basis of comparison was used.

## SECTION 2.0

### DESCRIPTION OF MODELS

A brief description of the several models that have been compared is presented here. Further details of this comparison can be found in Ref. 4 and in the original publications of each author. Most of the models include the effect of clouds, but this aspect of the models is not included here. Comparisons of cloud-cover global insolation models will follow (in a subsequent report) the clear sky model comparisons reported here. Such comparisons have also been performed by Davies and Hay [5].

#### 2.1 ATWATER AND BALL MODEL

Direct and global insolation models were published by Atwater and Ball [6,7]. The direct insolation model was taken from Kastrov as discussed by Kondratyev [8]. The equations of transfer and the transmission functions for this insolation model are given in Table 2-1. (The symbols in Tables 2-1 through 2-6 are defined in the Nomenclature.)

**Table 2-1. EQUATIONS FOR TOTAL DOWNWARD IRRADIANCE FOR ATWATER AND BALL MODEL**

---

#### Basic Equations

$$I_d = I_o (\cos Z) (T_{Md} - a_w) T_A$$

$$I_T = I_o (\cos Z) (T_M - a_w) T_A / (1 - r_g r_s)$$

#### Transmission Functions

$$T_{Md} = 1.041 - 0.16 [M(949 \times 10^{-6} P + 0.051)]^{0.5}$$

$$T_M = 1.021 - 0.0824 [M(949 \times 10^{-6} P + 0.051)]^{0.5}$$

$$a_w = 0.077 (U_w M)^{0.3}$$

$$T_A = \exp(-\tau_A M')$$

$$M = 35 / [(1224 \cos^2 Z) + 1]^{0.5}$$

$$M' = PM / 1013$$


---

The form of the equation for water vapor absorption was published by McDonald [9]. The value of  $r_s = 0.0685$  for a molecular atmosphere, as reported by Lacis and Hansen [10], was used with this model. Atwater and Ball

used MIE calculations to obtain  $\tau_A$ , which is much too rigorous for a simple model. Therefore, a value of  $\tau_A$  that will be described later in the Bird model was used here.

## 2.2 DAVIES AND HAY MODEL

A model for solar insolation (direct and diffuse) was published by Davies and Hay [5]. The equations used in this model were partially the result of comparing several existing models, and they are presented in Table 2-2.

**Table 2-2. EQUATIONS FOR TOTAL DOWNWARD IRRADIANCE FOR DAVIES AND HAY MODEL**

### Basic Equations

$$I_d = I_o (\cos Z) (T_o T_R - a_w) T_A$$

$$I_{as} = I_o (\cos Z) [T_o (1 - T_R) T_A (0.5) + (T_o T_R - a_w) (1 - T_A) W_o B_a]$$

$$I_G = r_g r_s (I_d + I_{as}) / (1 - r_g r_s)$$

$$I_T = I_d + I_{as} + I_G$$

### Transmission Functions

$$T_o = 1 - 0.02118X_o / (1 + 0.042X_o + 0.000323X_o^2) - 1.082X_o / (1 + 138.6X_o)^{0.805} - 0.0658X_o / [1 + (103.6X_o)^3]$$

$$X_o = U_o M$$

$$a_w = 2.9X_w / [(1 + 141.5X_w)^{0.635} + 5.925X_w]$$

$$X_w = U_w M$$

$$T_A = K^M$$

$$r_s = 0.0685 + (1 - B_a)(1 - T_A)W_o$$

The expressions for ozone transmittance,  $T_o$ , and water vapor absorption,  $a_w$ , were taken from Lacis and Hansen [10]. The transmission due to Rayleigh scattering,  $T_R$ , was presented in tabular form, and so we used the Bird model expression for  $T_R$  in this model. The value  $K = 0.91$  was used for data generated here and is representative of aerosol conditions in southern Ontario.  $W_o = 0.98$  and  $B_a = 0.85$  were used here also.

## 2.3 WATT MODEL

Another direct and diffuse insolation model has been constructed by Watt [3], based partially on the work of Moon [1]. The equations for this model are shown in Table 2-3.

**Table 2-3. EQUATIONS FOR TOTAL DOWNWARD IRRADIANCE FOR WATT MODEL**

### Basic Equations

$$I_d = I_o (\cos Z) T_{wa} T_{as} T_o T_{ws} T_L T_U$$

$$I_s = I_o [0.8 r_s (1 + r_g r_s) (1 + \cos Z)^{0.5} + 0.5 \alpha_{cs} r_g r_s \cos Z + 0.5 r_s \cos Z]$$

$$I_T = I_d + I_s$$

### Transmission Functions

$$T_w = 0.93 - 0.033 \log (U_w M_2)$$

$$T_{as} = 10^{-0.045 [(P/P_o) M_1]^{0.7}}$$

$$T_o = 10^{-(0.0071 + 0.01 U_o M_4)}$$

$$T_{ws} = 10^{-(0.0095 U_w M_2)}$$

$$T_L = 10^{\tau_L M_2^{0.7}}$$

$$T_U = 10^{-\tau_u M_3}$$

$$\tau_L = 0.6 (\tau_{0.5} - 0.01 U_w - 0.03)$$

$$\alpha_{cs} = (0.93 - 0.033 \log U_w) 10^{-[0.006 P/1013 + 0.4 (T_L + T_U)]}$$

$$r_s = \alpha_{cs} \{1 - 10^{-[0.003 P/1013 + 0.01 U_w + 0.4 (T_L + T_U)]}\}$$

$$M_i = \sec Z \text{ for } Z \leq 70^\circ \text{ (i = 1, 2, 3, 4)}$$

$$M_i = (h_2 F_{z2} - h_1 F_{z1}) / (h_2 - h_1) \text{ for } Z > 70^\circ \text{ (i = 1, 2, 3, 4)}$$

$$F_{zj} = \{ [(r/h_j) \cos Z]^2 + 2 r/h_j + 1 \}^{0.5} - (r/h_j) \cos Z \text{ (j = 1, 2)}$$

$$\text{for } i = 1, h_1 = 0 \text{ km and } h_2 = 30 \text{ km}$$

$$i = 2, h_1 = 0 \text{ km and } h_2 = 3 \text{ km}$$

$$i = 3, h_1 = 15 \text{ km and } h_2 = 25 \text{ km}$$

$$i = 4, h_1 = 20 \text{ km and } h_2 = 40 \text{ km}$$

The upper layer broadband turbidity,  $\tau_u$ , was not well defined by Watt. A value of  $\tau_u = 0.02$  was used in the calculations performed here, which appears to be an average value for locations in the United States. The parameter  $r$  is the earth's radius ( $6.4 \times 10^6$  m).

## 2.4 HOYT MODEL

The equations used in the model by Hoyt [11] are shown in Table 2-4.

Table 2-4. EQUATIONS FOR TOTAL DOWNWARD IRRADIANCE FOR HOYT MODEL

### Basic Equations

$$I_d = I_o (\cos Z) \left( 1 - \sum_{i=1}^5 a_i \right) T_{AS} T_R$$

$$I_{as} = I_o (\cos Z) \left( 1 - \sum_{i=1}^5 a_i \right) [(1 - T_R)0.5 + (1 - T_{AS})0.75]$$

$$I_G = (I_d + I_{as}) r_g \left( 1 - \sum_{i=1}^5 a_i' \right) [(1 - T_R')0.5 + (1 - T_{AS}')0.25]$$

$$I_T = I_d + I_{as} + I_G$$

### Transmission Functions

$$a_1 = a_w = 0.110 (0.75 U_w M + 6.31 \times 10^{-4})^{0.3} - 0.0121$$

$$a_2 = a_{co} = 0.00235 (126 M' + 0.0129)^{0.26} - 7.5 \times 10^{-4}$$

$$a_3 = (1 - T_o) = 0.045 (U_o M + 8.34 \times 10^{-4})^{0.38} - 3.1 \times 10^{-3}$$

$$a_4 = a_{ox} = 7.5 \times 10^{-3} (M')^{0.875}$$

$$a_5 = 0.05 T_{AS}$$

$$M' = MP/1013.25$$

Hoyt obtained air mass values,  $M$ , from Bemporad's [12] tables. The expression for air mass of Kasten [13] was used here instead. The values of  $T_{AS}$  and  $T_R$  are calculated from tables furnished by Hoyt [11]. The  $a_i'$  values are calculated using air mass values of  $M' + 1.66 P/1013.25$  in the  $a_i$  expressions.

$T_{AS}'$  and  $T_R'$  are evaluated for air mass values of 1.66 P/1013.25 in  $T_{AS}$  and  $T_R$ . The table values from which  $T_{AS}$  is calculated are limited so that large optical depths cannot be considered. Large optical depths can occur from high turbidity or from large zenith angles. In the data presented later for  $Z = 80^\circ$ , an approximate value of  $T_{AS}$  was used to complete the plotted results.

## 2.5 LACIS AND HANSEN MODEL

The equations for the model developed by Lacis and Hansen [10] are shown in Table 2-5.

**Table 2-5. EQUATIONS FOR TOTAL DOWNWARD IRRADIANCE FOR LACIS AND HANSEN MODEL**

### Basic Equation

$$I_T = I_0 (\cos Z) [(0.647 - r_s' - a_o)/(1 - 0.0685 r_g) + 0.353 - a_w]$$

### Transmission Functions

$$r_s' = 0.28/(1 + 6.43 \cos Z)$$

$$a_o = 1 - T_o \text{ as shown in Table 2-2}$$

$$a_w = \text{Shown in Table 2-2 with the following correction:}$$

$$X_w = X_w(P/1013)^{0.75}(273/T)^{0.5}$$

## 2.6 BIRD MODEL

A model has been constructed that is based on comparisons with the SOLTRAN 3 and SOLTRAN 4 [4] direct insolation models and the BRITE Monte Carlo global model [14]. Formalisms in the previous models that were considered to be optimum were adopted here. The equations for this model are shown in Table 2-6.

The atmospheric turbidity values,  $\tau_{A,0.38}$  and  $\tau_{A,0.5}$ , have been measured on a regular basis by the National Weather Service [15] at 0.38- and 0.5- $\mu\text{m}$  wavelengths, respectively. If one of the turbidity values is not available, its value can be entered as a zero in the expression for  $\tau_A$ . The expression for  $\tau_A$  is based on the Air Force Geophysics Laboratory (AFGL) rural aerosol model [16]. The expression used here for  $T_{AA}$  was found by fitting the expression to the results of the SOLTRAN 4 [4] code. The value of  $K_1 = 0.0933$  for the rural aerosol was used here for all calculations. For the urban aerosol model, which contains more carbon, the value of  $K_1 = 0.385$  was found to be appropriate. From a theoretical standpoint,  $K_1$  should be nearly equal to  $1 - W_o$ , where  $W_o$  is the single scattering albedo. The forward-scattering

**Table 2-6. EQUATIONS FOR TOTAL DOWNWARD IRRADIANCE FOR THE BIRD MODEL**
Basic Equations

$$I_d = I_o (\cos Z) (0.9662) T_R T_o T_{UM} T_w T_A$$

$$I_{as} = I_o (\cos Z) (0.79) T_o T_w T_{UM} T_{AA} \\ [0.5 (1 - T_R) + B_a (1 - T_{AS})] / [1 - M + (M)^{1.02}]$$

$$I_T = (I_d + I_{as}) / (1 - r_g r_s)$$

Transmission Equations

$$T_R = \exp \{-0.0903 (M')^{0.84} [1 + M' - (M')^{1.01}]\}$$

$$T_o = 1 - 0.1611 X_o (1 + 139.48 X_o)^{-0.3035} \\ - 0.002715 X_o (1 + 0.044 X_o + 0.0003 X_o^2)^{-1}$$

$$X_o = U_o M$$

$$T_{UM} = \exp [-0.0127 (M')^{0.26}]$$

$$T_w = 1 - 2.4959 X_w [(1 + 79.034 X_w)^{0.6828} + 6.385 X_w]^{-1}$$

$$X_w = U_w M$$

$$T_A = \exp [-\tau_A^{0.873} (1 + \tau_A - \tau_A^{0.7088}) M^{0.9108}]$$

$$\tau_A = 0.2758 \tau_{A,0.38} + 0.35 \tau_{A,0.5}$$

$$T_{AA} = 1 - K_1 (1 - M + M^{1.06}) (1 - T_A)$$

$$T_{AS} = T_A / T_{AA}$$

$$r_s = 0.0685 + (1 - B_a) (1.0 - T_{as})$$

$$M = [\cos Z + 0.15(93.885 - Z)^{-1.25}]^{-1}$$

$$M' = MP/1013$$

ratio,  $B_a$ , is related through MIE theory to a parameter  $\langle \cos \theta \rangle$ , called the asymmetry factor, by

$$B_a = 0.5(1 + \langle \cos \theta \rangle)$$

The asymmetry factor is the mean of the cosine of the scattering angle,  $\theta$ , with the angular intensity as the weighting function. The extreme values of  $B_a$  are

$$B_a = \begin{cases} 1 & \text{for all forward scattering;} \\ 0.5 & \text{for isotropic scattering;} \\ 0 & \text{for all backward scattering.} \end{cases}$$

Table 2-7 contains values of the asymmetry factor at various wavelengths for the rural aerosol model and the haze L aerosol model used by Dave [17]. Two differences in these aerosol models are: (1) the rural aerosol model is bimodal, whereas Dave's model has a single mode; and (2) the rural aerosol model varies the complex index of refraction with wavelength, and the Dave model holds it constant. The values of the asymmetry factors for the two models are in reasonably good agreement, and our calculations indicate that the Bird model is relatively insensitive to small changes in this parameter. A value of  $B_a = 0.82$  was used for the rural aerosol, and  $B_a = 0.86$  for Dave's aerosol in calculations shown later.

**Table 2-7. VALUES OF THE ASYMMETRY FACTOR FOR THE RURAL AND THE DAVE HAZE L AEROSOL**

Rural		Dave Haze L	
$\lambda$	$\langle \cos \theta \rangle$	$\lambda$	$\langle \cos \theta \rangle$
0.325	0.66	--	--
0.35	0.66	0.35	0.73
0.4	0.65	--	--
0.5	0.64	0.455	0.72
0.63	0.64	0.635	0.71
0.7525	0.63	0.7525	0.71
0.86	0.63	--	--
0.9935	0.63	0.994	0.70
1.235	0.64	1.235	0.69
1.497	0.65	1.61	0.67
1.8	0.68	2.1	0.63
2.198	0.71	2.198	0.62

It is suggested that values of  $B_a = 0.84$  and  $K_1 = 0.1$  be used with this model unless good information on the aerosol is available. All other data required by the model comes from meteorological measurements near the site of interest.

Ozone measurements are sometimes difficult to obtain. Since ozone has a minor effect on broadband solar insolation, it is suggested that the method of Van Heuklon [18] could be used in lieu of real data.

The expression used for air mass in the Bird model comes from Kasten [13] and was used for all of the calculations reported here except for the Atwater and Ball model and the Watt model.

For the convenience of the reader, Table 2-8 itemizes the input parameters required for each of the simple models.

**Table 2-8. INPUT PARAMETERS REQUIRED FOR SIMPLE MODELS**

Model	Input
Atwater and Ball	Solar constant, zenith angle, surface pressure, ground albedo, precipitable water vapor, total ozone, broadband turbidity
Davies and Hay	Solar constant, zenith angle, surface pressure, ground albedo, precipitable water vapor, total ozone, aerosol single scattering albedo (0.98 suggested), aerosol forward scattering ratio (0.85 suggested), broadband aerosol transmittance
Watt	Solar constant, zenith angle, surface pressure, ground albedo, precipitable water vapor, total ozone, turbidity at 0.5- $\mu$ m wavelength, upper layer turbidity
Hoyt	Solar constant, zenith angle, surface pressure, ground albedo, precipitable water vapor, total ozone, turbidity at one wavelength
Lacis and Hansen	Solar constant, zenith angle, surface pressure, surface temperature, ground albedo, precipitable water vapor, total ozone
Bird	Solar constant, zenith angle, surface pressure, ground albedo, precipitable water vapor, total ozone, turbidity at 0.5- and/or 0.38- $\mu$ m wavelength, aerosol forward scattering ratio (0.84 recommended)

## 2.7 RIGOROUS CODES

Three rigorous codes have been used in this study as a basis for formulating the Bird model. One code is for direct normal irradiance and is called SOLTRAN 4 [4]. Two other codes, which include both the direct normal and the diffuse irradiance, are the BRITE [14] Monte Carlo code and the Dave [17] Spherical Harmonics code. In each of these codes, a multilayered atmospheric model can be constructed, where important atmospheric parameters are defined

at each layer. In this way, a fairly detailed atmosphere can be constructed that closely resembles the real atmosphere at a given time and specific location. Each code then uses its own technique to solve the radiative transfer problem.

These rigorous codes calculate the irradiance at a specified altitude, sun angle, and for an atmospheric model at discrete wavelengths. For comparison with the simple broadband models described previously, the spectral irradiance from the rigorous codes has to be integrated over wavelength.

The SOLTRAN 3 code, an earlier version of SOLTRAN 4, was used to formulate most of the transmittance functions found in the Bird model. This was accomplished by performing a least-square fit of each transmittance function to transmittance data from SOLTRAN 3 for each atmospheric constituent. Further details of this operation can be found in Ref. 4. SOLTRAN 3 was used because it was the only version available to us when this portion of the work was performed. The only difference in the two models that would be apparent in the results would be caused by a slight difference in the "continental" and "rural" aerosol models that are used in the two codes.

A comparison was made between data from the BRITE code and several of the simple clear sky global models. Based on this comparison and the author's judgment of the best expressions used in the simple models, a model for the diffuse irradiance was formulated. This simple model of the diffuse irradiance was then fine-tuned to provide good agreement with the BRITE code as well as results from the Dave Spherical Harmonics code.

It is appropriate to comment here that there are problems with the expressions used for the diffuse irradiance model. The general formalism for the diffuse transfer equation of some of the simple models was adopted even though it may not be as acceptable, based on the physics of the problem, as one would like. For example, a cosine of the solar zenith angle is included in the diffuse transfer equation. This implies that all of the diffuse radiation behaves just like the direct normal component. The cosine is used to calculate the irradiance falling on a horizontal surface. However, it is well known that the diffuse irradiance is much more complex than this. An example of a more rigorous but fairly simple formalism for the diffuse irradiance is found in Ref. 19, in which the diffuse radiation is divided into three components: an isotropic term, a term resulting from horizon brightening, and a circumsolar term. The circumsolar term is the only one that behaves very much like the direct normal radiation. For tilted surfaces, a ground reflection should be added to this diffuse model. Another problem with this formalism is associated with using transmittance expressions for diffuse radiation that were derived for direct radiation.

## SECTION 3.0

## MODEL COMPARISONS

Each of the simple models described in Section 2.0 was programmed on a computer to produce data for comparison. A comparison of the aerosol transmittance, the transmittance after molecular (Rayleigh) scattering, the water vapor ( $H_2O$ ) transmittance, and the ozone ( $O_3$ ) transmittance will be presented first. Then, a comparison between the direct, the diffuse sky, the diffuse sky/ground, and the global radiation for three different atmospheric models will be shown. Comparisons are made, where possible, between each of the simple model results as well as the results from the rigorous models.


## 3.1 TRANSMITTANCE COMPARISONS

To become oriented as to the relative importance of each atmospheric constituent in atmospheric transmittance, the broadband transmittance versus the secant of the solar zenith angle (approximate air mass) for each constituent was plotted in Fig. 3-1. This figure was generated with output from the SOLTRAN 3 code for a Midlatitude Summer (MLS) atmospheric model. Table 3-1 shows the amounts of  $H_2O$  and  $O_3$  from sea level to the top of the atmosphere in a vertical column for the two atmospheric models--MLS and USS (U.S. Standard)--used in this comparison.

Table 3-1. AMOUNTS OF  $H_2O$  AND  $O_3$  IN A VERTICAL COLUMN FOR THE MLS AND USS ATMOSPHERES

	$H_2O$ (cm)	$O_3$ (cm)
MLS	2.93	0.31
USS	1.42	0.34

An examination of Fig. 3-1 shows that  $CO_2$  and  $O_2$  are the least important attenuators, and this is why they are not included in some models. The next element exhibiting increased attenuation is  $O_3$ , followed by  $H_2O$ . Molecular scattering dominates the total molecular absorption at large zenith angles and has a greater effect than most individual molecular species at all zenith angles. The most significant attenuator at all zenith angles is the aerosol. The aerosol modeled here was the continental aerosol model [16] with a sea level meteorological range of 23 km. A 23-km meteorological range is considered to be a moderately clear atmosphere. This aerosol produces an optical depth of 0.27 (base e) at 0.5- $\mu m$  wavelength.



**Figure 3-1. Transmittance versus Secant of Solar Zenith Angle for Midlatitude Summer Model**

The data in Fig. 3-1 indicate the relative effect that the atmospheric constituents have on the direct irradiance. However, when global irradiance is being considered these results can be misleading. A large fraction of the radiation lost in the direct beam by molecular and aerosol scattering is regained in the diffuse component. As a result of this, changes in aerosol optical depth through the atmosphere have less effect on the total global irradiance than on the direct irradiance.

Figures 3-2 and 3-5 present transmittance data for the USS atmosphere from several of the models. It should be pointed out that the broadband turbidity expression from the Bird model was used in the Atwater and Ball model. A similar effort could have been made with the aerosol transmittance term in the Davies and Hay model to produce identical results. One of the strengths of the Bird model is that it is based entirely on algebraic expressions for transmittance calculations rather than tabulated data. This makes the use of the model considerably easier and provides more flexibility.

The comparison made here for one model atmosphere is not really indicative of the accuracy of each model. Since a wide range of values of turbidity,  $H_2O$  amount, and  $O_3$  amount are required for real atmospheric conditions, the model must be able to accommodate these changes. Additional comparisons are presented in Ref. 4 for a range of these parameters. As was stated earlier, the transmittance expressions in the Bird model were derived from comparisons with SOLTRAN 3 results, but the comparisons made here are with SOLTRAN 4 results. This should have an effect only on the aerosol transmittance shown in Fig. 3-2. The transmittance comparisons shown in Figs. 3-2 through 3-5 are rather self-explanatory, and so no further discussions are presented.

### 3.2 IRRADIANCE COMPARISONS

The global solar irradiance has been divided into three components: the direct irradiance on a horizontal surface, the diffuse sky irradiance on a horizontal surface, and the diffuse ground/sky irradiance on a horizontal surface. The diffuse sky irradiance is the total diffuse radiation present when the ground has zero albedo (completely absorbing ground), and the diffuse ground/sky irradiance is that amount added to the total diffuse irradiance when the ground albedo is not zero.

Figures 3-6 through 3-9 present comparisons of the global irradiance at sea level in the USS atmosphere from all of the models as well as comparisons of the three components of the global irradiance. For this atmospheric model, the Bird, Hoyt, and Monte Carlo models produce very similar results. The model of Atwater and Ball significantly underestimates the diffuse sky irradiance, and the Watt model overestimates the diffuse sky irradiance. The results of the Davies and Hay model would have shown much better agreement with the Monte Carlo results if a more reasonable value of the aerosol transmittance had been used. The Bird model Rayleigh transmittance was used in the Davies and Hay model.

It is instructive to examine the relative magnitude of each component. For a solar zenith angle of zero (the sun directly overhead), the direct component provides approximately 81% of the total, the diffuse sky approximately 17%,




Figure 3-2. Aerosol Transmittance - USS Atmosphere (23-km-Visibility Aerosol)







Figure 3-3. Rayleigh Transmittance




**Figure 3-4. Water Vapor Transmittance – USS Atmosphere  
(1.42-cm H<sub>2</sub>O in Vertical Column)**




**Figure 3-5. Ozone Transmittance – USS Atmosphere**




**Figure 3-6. Total Insolation – USS Atmosphere (23-km-Visibility Rural Aerosol, Albedo = 0.2)**



**Figure 3-7. Direct Horizontal Insolation – USS Atmosphere (V = 23 km; U<sub>w</sub> = 1.42 cm; U<sub>o</sub> = 0.34 cm)**



**Figure 3-8. Diffuse Sky Insolation – USS Atmosphere ( $V = 23$  km;  $U_w = 1.42$  cm;  $U_o = 0.34$  cm)**



**Figure 3-9. Diffuse Ground/Sky Insolation – USS Atmosphere (23-km-Visibility Rural Aerosol, Albedo = 0.2)**

and the diffuse ground/sky approximately 2%. The calculations are for a turbidity of 0.27 at 0.5- $\mu\text{m}$  wavelength and a ground albedo of 0.2.


In Figures 3-10 through 3-13, a similar set of plots is presented for the MLS atmospheric model with an atmospheric turbidity of 0.27 at 0.5- $\mu\text{m}$  wavelength and a ground albedo of 0.8. The larger ground albedo increases the diffuse ground/sky component by a factor of four. It is evident that the simple models begin to deviate from the Monte Carlo result for the diffuse ground/sky component. However, since this component is so small, the global result is still in close agreement. The relative agreement of the results from the different models is nearly the same as with the USS atmosphere. This is principally because the aerosol model is identical in both models.

Finally, a comparison is made of the models for the MLS atmosphere with the Haze L aerosol model of Dave [17]. The Dave atmosphere modeled here consists of 15 homogeneous layers instead of the 32 exponentially varying layers that were used in the previous MLS atmosphere. In addition, the Haze L aerosol model is significantly different from the rural aerosol model used previously. Not only are the particle size distributions and complex indices of refraction different, but most importantly the number density of the aerosol as a function of altitude is very different. The turbidity of this model is 0.0996 instead of the 0.266 used previously in the vicinity of 0.5- $\mu\text{m}$  wavelength. Calculations with SOLTRAN 4 show that a turbidity of 0.0996 in the rural aerosol model corresponds to a sea level visibility of nearly 250 km. This is an extremely clear atmosphere. Figures 3-14 through 3-17 illustrate the comparison results for this atmospheric model.


It is readily apparent from the results shown in Figs. 3-14 through 3-17 that the Atwater and Ball model is based on a very clear atmosphere, since it agrees much better with the Dave data. Similarly, the aerosol parameter,  $K = 0.91$ , used in the Davies and Hay model is for a very clear atmosphere. The Lacis and Hansen model appears to be in slightly closer agreement with this clear atmosphere also, but it does not have provisions for changes in turbidity.

The clear sky diffuse irradiance of the Atwater and Ball model as shown in the figures presented here may be slightly lower than the model intended because of the way the calculations were performed. This model is really composed of two separate models: one for the direct irradiance and one for the global irradiance. The clear sky diffuse irradiance was obtained by running the model for a ground albedo of zero and then subtracting the direct horizontal from the global. If the direct horizontal irradiance is slightly high, as it appears to be, then the diffuse term would be slightly lower than expected. The real evaluation of this model should be made on the global horizontal irradiance.


It will be noted in Fig. 3-14 that there is a slight difference between the Monte Carlo global and the spherical harmonic global results of Dave ( $\sim 3.6\%$  at a zenith angle of 0). Figure 3-15 shows that a large fraction of this difference is in the direct component. Since the direct component of the Monte Carlo code is deterministic in nature rather than statistical, these differences are most likely due to differences in the molecular absorption coefficients used and the band absorption models used. Dave used an older set




**Figure 3-10. Total Insolation – MLS Atmosphere (23-km-Visibility Rural Aerosol, Albedo = 0.8)**




**Figure 3-11. Direct Horizontal Insolation – MLS Atmosphere ( $V = 23$  km,  $U_w = 2.93$  cm,  $U_o = 0.31$  cm)**




**Figure 3-12. Diffuse Sky Insolation — MLS Atmosphere ( $V = 23$  km,  $U_w = 2.93$  cm,  $U_o = 0.31$  cm)**




**Figure 3-13. Diffuse Ground/Sky Insolation — MLS Atmosphere (23-km-Visibility Rural Aerosol, Albedo = 0.8)**




**Figure 3-14. Total Insolation—Dave Model 3 (Albedo = 0.2)**



**Figure 3-15. Direct Horizontal Insolation—Dave Model 3 ( $U_w = 2.93$  cm,  $U_o = 0.31$  cm)**




**Figure 3-16. Diffuse Sky Insolation—Dave Model 3**  
 $(U_w = 2.93 \text{ cm}, U_o = 0.31 \text{ cm})$




**Figure 3-17. Diffuse Ground/Sky Insolation—Dave Model 3**  
 $(\text{Albedo} = 0.2)$

of molecular absorption coefficients from AFGL than was used in the Monte Carlo code [20]. Figure 3-18 is a comparison of the spectral direct irradiance for these two codes. It is evident that there are some weak absorption bands present in the Monte Carlo code that are not present in the spherical harmonics code, and the shape of some of the bands is significantly different for the two codes. SOLTRAN 4 is based on the same absorption data that Dave used, and similar differences in the direct normal irradiance occur between SOLTRAN 4 and the Monte Carlo code. These differences are shown in Fig. 3-19, which has 31 more data points in the Monte Carlo results than in Fig. 3-18. This increase in the number of data points increases the apparent spectral resolution. The SOLTRAN 4 code provided approximately 600 data points in this figure. Our conclusion is that most of the differences in the results from the rigorous codes are due to differences in the molecular absorption coefficients used.

A final observation is that many of the simple models have been based on actual measured data rather than comparison to rigorous models. This fact can make a difference in the direct normal irradiance or the diffuse irradiance but should not affect the total irradiance. The reason for this is that pyrheliometers measure the irradiance in a 5.8-degree field-of-view, which includes some diffuse or circumsolar irradiance. The rigorous codes include only the direct normal irradiance with no circumsolar. This means that the direct normal irradiance calculated with the Bird model will slightly underestimate the irradiance measured by a pyrheliometer. On a normal clear day, one is talking about less than a 1% underestimation. Let us reiterate that the total insolation should agree.



**Figure 3-18. Direct Normal Insolation — Dave Model 3 (AM1)**



**Figure 3-19. Direct Normal Insolation—USS Atmosphere (23-km-Visibility Rural Aerosol, AM1)**

## SECTION 4.0

### SUMMARY AND CONCLUSIONS

Five simple broadband models for clear sky global horizontal insolation have been compared with the spectrally integrated results from three rigorous spectral codes. As a part of this comparison, a sixth simple broadband model has been formulated. This sixth model, designated the Bird model, uses parts of the formalisms from the other simple models and has been fine-tuned to provide good agreement with the rigorous codes. The Bird model was constructed so that readily available meteorological data could be used in it. It is based entirely on algebraic expressions rather than look-up tables, which greatly simplifies the use of the model.

The comparison of the results from each of the simple models with the results of the rigorous codes indicates the following:

- The Atwater and Ball model is applicable to extremely clear atmospheric conditions with an atmospheric turbidity (base e) near 0.1- at 0.5- $\mu\text{m}$  wavelength. For turbidities near 0.27, this model underestimated the global irradiance by approximately 8% for air mass 1 (AM1). This model is extremely simple but does not have a good method of treating aerosol transmittance.
- The Watt model is relatively complicated and appears to overestimate the global insolation for AM1 conditions by approximately 7%. This is a complete model based on meteorological parameters. However, the upper air turbidity required in this model is not readily available.
- The Hoyt model provides excellent agreement with the rigorous codes. However, its use of look-up tables and the requirement to recalculate transmittance and absorptance parameters for modified air mass values causes this model to be relatively difficult to use.
- The Lacis and Hansen model is extremely simple. It tends to overestimate the global irradiance by approximately 8% at AM1, and it has no provisions for calculating direct irradiance.
- The Davies and Hay model could possibly provide good agreement with the rigorous codes. However, it uses a look-up table for the Rayleigh scattering transmittance term and does not have a good method for treating aerosol transmittance. The aerosol transmittance through a vertical path used by Davies and Hay for southern Ontario ( $K = 0.91$ ) is for an extremely clear atmosphere.
- It is hoped that the rigorous codes and accurate simple models will provide results that will agree within  $\pm 5\%$  with quality experimental data on clear days. Cloudy days are much more difficult to model accurately, and clouds can have the greatest effect on the total irradiance. Models that address cloud influences for irradiance will be examined at a later date.

- It should be recalled that the basis of comparison/evaluation of the simple models is the much more rigorous radiative transfer codes--as opposed to a comparison with actual data. Because of a lack of suitable, high-quality data, comparisons with actual data are impossible at this time. The greatest deficiency has been the lack of meteorological measurements accompanying good insolation data. However, efforts\* are currently underway at SERI and several universities to provide such data. As this data becomes available, comparisons and improvements will be made. Until then, it appears that both the Hoyt and Bird simplified models yield results in good agreement with the rigorous techniques. However, the Bird model is more flexible and easily used.

---

\*As part of the U.S. Department of Energy's Insolation Resource Assessment Program.

## SECTION 5.0

## REFERENCES

1. Moon, P. "Proposed Standard Solar-Radiation Curves for Engineering Use." J. Franklin Institute. Vol. 230: pp. 583-617; 1940.
2. Mahaptra, A. K. "An Evaluation of a Spectroradiometer for the Visible-Ultraviolet and Near-Ultraviolet." Ph.D. dissertation; University of Missouri; Columbia, MO; 1973; 121 pp. (University Microfilms 74-9964).
3. Watt, D. On the Nature and Distribution of Solar Radiation. HCP/T2552-01. U.S. Department of Energy; 1978.
4. Bird, R. E.; Hulstrom, R. E. Direct Insolation Models. SERI/TR-335-344. Golden, CO: Solar Energy Research Institute; 1980.
5. Davies, J. A.; Hay, J. E. "Calculation of the Solar Radiation Incident on a Horizontal Surface." Proceedings, First Canadian Solar Radiation Data Workshop. April 17-19, 1978. Canadian Atmospheric Environment Service; 1979.
6. Atwater, M. A.; Ball, J. T. "A Numerical Solar Radiation Model Based on Standard Meteorological Observations." Solar Energy. Vol. 21: pp. 163-170; 1978.
7. Atwater, M. A.; Ball, J. T. Solar Energy. Vol. 23: p. 275; 1979.
8. Kondratyev, K. Ya. Radiation in the Atmosphere. New York: Academic Press; 1969.
9. McDonald, J. E. "Direct Absorption of Solar Radiation by Atmospheric Water Vapor." J. Meteorology. Vol. 17: pp. 319-328; 1960.
10. Lacis, A. L.; Hansen, J. E. "A Parameterization for Absorption of Solar Radiation in the Earth's Atmosphere." J. Atmospheric Science. Vol. 31: pp. 118-133; 1974.
11. Hoyt, D. V. "A Model for the Calculation of Solar Global Insolation." Solar Energy. Vol. 21: pp. 27-35; 1978.
12. Bemporad, A. Zur Theorie der Extinktion des Lichtes in der Erdatmosphäre. Mitteilungen der Grossherzoglichen Sternwarte Zu Heidelberg; No. 4; 1904.
13. Kasten, F. A New Table and Approximation Formula for the Relative Optical Air Mass. Technical Report 136. Hanover, New Hampshire: U.S. Army Material Command, Cold Region Research and Engineering Laboratory; 1964.
14. Bird, R. E.; Hulstrom, R. L. Application of Monte Carlo Techniques to Insolation Characterization and Prediction. SERI/RR-36-306. Solar Energy Research Institute; Golden, CO; 1979.

15. Flowers, E. C.; McCormick, G. A.; Kurfis, K. R. "Atmospheric Turbidity over the United States." J. Appl. Meteor. Vol. 8: pp. 955-962; 1969.
16. Shettle, E. P.; Fenn, R. W. "Models of the Atmospheric Aerosol and Their Optical Properties." Proceedings of AGARD Conference No. 183, Optical Propagation in the Atmosphere. pp. 2.1-2.16; Presented at the Electromagnetic Wave Propagation Panel Symposium, Lyngby, Denmark; 27-31 October 1975.
17. Dave, J. V. "Extensive Datasets of the Diffuse Radiation in Realistic Atmospheric Models with Aerosols and Common Absorbing Gases." Solar Energy. Vol. 21: pp. 361-369; 1978.
18. Van Heuklon, T. K. "Estimating Atmospheric Ozone for Solar Radiation Models." Solar Energy. Vol. 22: pp. 63-68; 1979.
19. Hooper, F. C.; Brunger, A. P. "A Model for the Angular Distribution of Sky Radiance." Presented at the Joint ASME/AIChE 18th National Heat Transfer Conference. San Diego, CA; 6-8 August 1979.
20. Rothman, L. S. "Update of AFGL Atmospheric Absorption Line Parameters Compilation." Applied Optics. Vol. 17: pp. 3517-3518; 1978.

**APPENDIX A**  
**TABULATED MODEL DATA**

Three sets of tabulated data from each simple model are presented here, and each set represents one of the atmospheric models discussed in the main text. The parameters listed at the top of each table are:

- $I_0$  = Solar constant ( $W/m^2$ )
- UW = Precipitable water vapor in vertical path (cm)
- UO = Ozone amount in vertical path (cm)
- PR = Surface pressure (mb)
- TAU5 = Turbidity at 0.5- $\mu m$  wavelength
- TAU38 = Turbidity at 0.38- $\mu m$  wavelength
- RS = Ground albedo
- TEMP = Surface temperature (K)
- CONST = Constant K used in BIRD model
- BA = Forward to total scattered irradiance ratio
- TAUB = Broadband turbidity

The parameters at the top of each column of data are self-explanatory for the transmittance and absorptance terms. The remaining parameters are:

- DIRH = Direct horizontal irradiance ( $W/m^2$ )
- DIFSH = Diffuse sky horizontal irradiance ( $W/m^2$ )
- DIFGH = Diffuse ground horizontal irradiance ( $W/m^2$ )
- DTOT = Total or global horizontal irradiance ( $W/m^2$ )

Table A-1. TABULATED DATA FROM SEVERAL MODELS FOR THE  
 USS ATMOSPHERE

IO = 1353.0000  
 UW = 1.4200  
 UO = 3400  
 PR = 1013.0000  
 TAUS = 2661  
 TAU38 = 3538  
 RS = 2000  
 TEMP = 288.1000  
 CONST = 0.933  
 BA = 8200  
 TAUB = 1.907

ZENITH	AIRMASS
0.0000	9995
20.0000	1.0634
30.0000	1.1536
48.1900	1.4972
50.0000	1.5525
60.0000	1.9927
70.0000	2.8997
75.0000	3.8076
80.0000	5.5790
85.0000	10.3163

## ATWATER MODEL

ZEN	TA	AW	TMD	DIRH	DIFSH	DIFGH	DTOT
0.0	.8264	.0855	.9381	888.2846	64.9349	13.2405	966.4600
20.0	.8163	.0872	.9355	817.6250	62.8348	12.2298	892.6897
30.0	.8024	.0893	.9319	732.1165	60.0704	11.0037	803.1906
48.2	.7513	.0966	.9195	506.4300	51.2334	7.7461	565.4095
50.0	.7434	.0977	.9176	480.1187	50.0130	7.3637	537.4954
60.0	.6832	.1053	.9038	327.3209	41.7589	5.1266	374.2064
70.0	.5736	.1179	.8795	172.0532	30.0717	2.8076	204.9324
75.0	.4806	.1281	.8585	100.5358	22.3903	1.7075	124.6335
80.0	.3382	.1441	.8233	40.7697	13.2023	.7497	54.7216
85.0	.1249	.1752	.7472	4.9225	3.5038	.1170	8.5433

## WATT MODEL

ZEN	TA	TH20A	TH20S	TO3	TAIR	DIRH	DIFSH	DIFGH	DTOT
0.0	.8653	.9250	.9694	.9761	.9016	923.885	176.483	11.272	1111.639
20.0	.8590	.9241	.9675	.9756	.8974	854.888	171.361	10.680	1036.929
30.0	.8503	.9229	.9648	.9749	.8917	771.205	165.065	9.957	946.227
48.2	.8187	.9192	.9545	.9723	.8714	548.932	147.788	7.996	704.717
50.0	.8138	.9186	.9528	.9719	.8683	522.831	145.694	7.761	676.286
60.0	.7770	.9150	.9398	.9685	.8451	369.969	133.040	6.352	509.361
70.0	.7092	.9096	.9132	.9615	.8029	210.444	118.758	4.789	333.991
75.0	.6544	.9056	.8873	.9562	.7702	135.596	111.100	3.963	250.659
80.0	.5620	.9000	.8373	.9456	.7142	67.205	103.151	3.117	173.473
85.0	.3961	.8904	.7074	.9259	.6123	16.682	94.951	2.255	113.888

## HOYT MODEL

ZEN	TAS	AA	AW	ACO2	A03	A02	TR
0.0	.8317	.0416	1000	.0075	.0268	.0075	.9170
20.0	.8220	.0411	1021	.0076	.0275	.0079	.9125
30.0	.8084	.0404	1049	.0078	.0285	.0085	.9063
48.2	.7588	.0379	1144	.0084	.0317	.0107	.8840
50.0	.7511	.0376	1158	.0085	.0322	.0110	.8807
60.0	.6926	.0346	1258	.0091	.0357	.0137	.8560
70.0	.5859	.0293	1422	.0101	.0417	.0190	.8138
75.0	.4956	.0248	1553	.0109	.0466	.0242	.7781

ZEN	DIRH	DIFSH	DIFGH	DTOT
0.0	842.7127	185.2787	19.8473	1047.8387
20.0	776.0145	183.3879	18.4825	977.8849
30.0	695.2566	180.8148	16.8257	892.8972
48.2	482.1123	171.6728	12.4167	666.2018
50.0	457.2772	170.2822	11.8978	639.4573
60.0	313.2352	159.8684	8.8496	481.9533
70.0	167.1730	141.5303	5.6278	314.3311
75.0	99.6955	126.4744	4.0258	230.1957

Table A-1. TABULATED DATA FROM SEVERAL MODELS FOR THE  
 USS ATMOSPHERE (concluded)

LACIS MODEL				
ZEN	AW	A03	DTOT	
0.0	.1089	.0234	1134.0234	
20.0	.1109	.0240	1059.6338	
30.0	.1135	.0249	969.0372	
48.2	.1220	.0283	725.7510	
50.0	.1233	.0288	676.8444	
60.0	.1319	.0327	525.3496	
70.0	.1455	.0403	339.5933	
75.0	.1559	.0474	244.5878	
80.0	.1711	.0602	150.9857	
85.0	.1969	.0906	63.4795	

DAVIES MODEL				
ZEN	DIRH	DIFSH	DIFGH	DTOT
0.0	964.6041	131.3193	18.2102	1114.1335
20.0	893.1822	129.3610	17.1608	1039.7040
30.0	806.4406	126.7305	15.8783	949.0494
48.2	575.4248	117.7340	12.3977	705.5564
50.0	548.2303	116.4077	11.9791	676.6171
60.0	388.6384	106.7716	9.4598	504.8698
70.0	221.5297	90.5812	6.6085	318.7193
75.0	140.8761	77.6703	5.0465	223.5929
80.0	68.5779	58.6164	3.3589	130.5533
85.0	16.1786	29.9684	1.5249	47.6719

BIRD MODEL							
ZEN	TA	T03	TU	TR	TAS	AW	TAA
0.0	.8127	.9822	.9874	.9137	.8271	.1032	.9825
20.0	.8029	.9814	.9872	.9094	.8180	.1048	.9815
30.0	.7895	.9803	.9869	.9033	.8055	.1068	.9801
48.2	.7410	.9763	.9860	.8816	.7603	.1134	.9746
50.0	.7336	.9756	.9859	.8783	.7541	.1144	.9728
60.0	.6778	.9709	.9849	.8531	.7029	.1209	.9643
70.0	.5785	.9619	.9834	.8074	.6126	.1312	.9444
75.0	.4959	.9537	.9822	.7684	.5435	.1389	.9124
80.0	.3704	.9391	.9803	.7078	.4494	.1499	.8242
85.0	.1757	.9056	.9770	.6157	.4256	.1684	.4130

ZEN	DIRH	DIFSH	DIFGH	DTOT
0.0	844.2037	168.9023	20.5954	1033.7014
20.0	777.9391	166.1341	19.5137	963.5869
30.0	697.7136	162.3802	18.1816	878.2754
48.2	485.8908	149.7404	14.5168	650.1480
50.0	461.1879	147.4221	14.0423	622.6523
60.0	317.6805	133.6742	11.2868	462.6415
70.0	171.6204	110.5682	8.0237	290.2122
75.0	103.9955	90.4421	6.0411	200.4786
80.0	46.5685	59.8652	3.6917	110.1254
85.0	9.0703	12.5752	.7707	22.4162

Table A-2. TABULATED DATA FROM SEVERAL MODELS FOR THE  
MLS ATMOSPHERE

ID = 1353.0000  
 UW = 2.9300  
 UO = 3100  
 PR = 1013.0000  
 TAU5 = 2661  
 TAU30 = 3538  
 RS = 8000  
 TEMP = 294.0000  
 CONST = .0933  
 BA = 8200  
 TAUB = 1907

ZENITH	AIRMASS
0.0000	.9995
20.0000	1.0634
30.0000	1.1536
48.1900	1.4972
50.0000	1.5525
60.0000	1.9927
70.0000	2.8997
75.0000	3.8076
80.0000	5.5790
85.0000	10.3163

ATWATER MODEL

ZEN	TA	AW	TMD	DIRH	DIFSH	DIFGH	DTOT
0.0	.8264	.1063	.9381	865.0702	64.9349	53.9190	983.9241
20.0	.8163	.1083	.9355	795.6702	62.8348	49.7737	908.2787
30.0	.8024	.1110	.9319	711.7362	60.0704	44.7471	816.5538
48.2	.7513	.1200	.9195	490.5418	51.2334	31.4106	573.1858
50.0	.7434	.1214	.9176	464.7946	50.0130	29.8471	544.6547
60.0	.6832	.1308	.9038	315.5108	41.7589	20.7135	377.9832
70.0	.5736	.1465	.8795	164.4570	30.0717	11.2782	205.8069
75.0	.4806	.1592	.8585	95.3030	22.3903	6.8235	124.5168
80.0	.3382	.1790	.8233	37.9910	13.2023	2.9680	54.1614
85.0	.1249	.2177	.7472	4.2961	3.5038	.4522	8.2521

WATT MODEL

ZEN	TA	TH20A	TH20S	T03	TAIR	DIRH	DIFSH	DIFGH	DTOT
0.0	.8836	.9146	.9379	.9768	.9016	903.091	220.790	58.727	1182.608
20.0	.8779	.9137	.9341	.9763	.8974	834.679	214.383	55.738	1104.800
30.0	.8701	.9125	.9287	.9757	.8917	751.725	206.506	52.083	1010.315
48.2	.8417	.9088	.9083	.9733	.8714	531.559	184.892	42.168	758.619
50.0	.8373	.9083	.9051	.9729	.8683	505.732	182.271	40.978	728.982
60.0	.8038	.9047	.8797	.9698	.8451	354.681	166.441	33.849	554.970
70.0	.7413	.8992	.8291	.9635	.8029	197.827	148.573	25.932	372.332
75.0	.6904	.8953	.7813	.9586	.7702	124.844	138.992	21.750	285.586
80.0	.6032	.8896	.6933	.9489	.7142	59.236	129.047	17.457	205.740
85.0	.4434	.8800	.4896	.9309	.6123	12.842	118.790	13.085	144.716

HOYT MODEL

ZEN	TAS	AA	AW	ACD2	A03	A02	TR
0.0	.8317	.0416	.1272	.0075	.0258	.0075	.9170
20.0	.8220	.0411	.1298	.0076	.0264	.0079	.9125
30.0	.8084	.0404	.1333	.0078	.0274	.0085	.9063
48.2	.7588	.0379	.1451	.0084	.0305	.0107	.8840
50.0	.7511	.0376	.1469	.0085	.0310	.0110	.8807
60.0	.6926	.0346	.1592	.0091	.0344	.0137	.8560
70.0	.5859	.0293	.1796	.0101	.0401	.0190	.8138
75.0	.4956	.0248	.1960	.0109	.0448	.0242	.7781

ZEN	DIRH	DIFSH	DIFGH	DTOT
0.0	815.7084	179.3415	72.9219	1067.9718
20.0	750.5957	177.3810	67.8237	995.8004
30.0	671.8142	174.7181	61.6398	908.1721
48.2	464.2596	165.3157	45.2155	674.7908
50.0	440.1194	163.8930	43.2865	647.2989
60.0	300.3540	153.2941	31.9775	485.6256
70.0	159.2519	134.8242	20.0839	314.1600
75.0	94.4394	119.8065	14.2080	228.4540

Table A-2. TABULATED DATA FROM SEVERAL MODELS FOR THE  
MLS ATMOSPHERE (concluded)

LACIS MODEL

ZEN	AW	A03	DTOT
0.0	.1327	.0224	1138.0919
20.0	.1350	.0231	1063.0223
30.0	.1379	.0239	971.6311
48.2	.1475	.0270	726.4310
50.0	.1489	.0275	697.3221
60.0	.1585	.0312	524.7679
70.0	.1736	.0382	338.2250
75.0	.1849	.0448	243.0373
80.0	.2012	.0568	149.4940
85.0	.2282	.0851	62.4267

DAVIES MODEL

ZEN	DIRH	DIFSH	DIFGH	DTOT
0.0	936.3331	129.0411	74.5253	1139.8995
20.0	866.5097	127.0680	70.2349	1063.8126
30.0	781.7525	124.4193	64.9933	971.1651
48.2	556.3066	115.3767	50.7790	722.4623
50.0	529.8009	114.0456	49.0708	692.9173
60.0	374.4382	104.3899	38.7949	517.6231
70.0	212.2479	88.2198	27.1739	327.6415
75.0	134.2798	75.3692	20.8053	230.4543
80.0	64.7464	56.4768	13.9067	135.1298
85.0	14.9323	28.3066	6.3442	49.5832

BIRD MODEL

ZEN	TA	T03	TU	TR	TAS	AW	TAA
0.0	.8127	.9834	.9874	.9137	.8271	.1219	.9825
20.0	.8029	.9826	.9872	.9094	.8180	.1235	.9815
30.0	.7895	.9816	.9869	.9033	.8055	.1257	.9801
48.2	.7410	.9778	.9860	.8816	.7603	.1330	.9746
50.0	.7336	.9772	.9859	.8783	.7541	.1340	.9728
60.0	.6778	.9727	.9849	.8531	.7029	.1411	.9643
70.0	.5785	.9643	.9834	.8074	.6126	.1520	.9444
75.0	.4959	.9566	.9822	.7684	.5435	.1601	.9124
80.0	.3704	.9430	.9803	.7078	.4494	.1717	.8242
85.0	.1757	.9116	.9770	.6157	.4256	.1907	.4130

ZEN	DIRH	DIFSH	DIFGH	DTOT
0.0	827.6234	165.5850	86.0091	1079.2176
20.0	762.5502	162.8477	81.5689	1006.9668
30.0	683.7850	159.1385	76.1007	919.0243
48.2	475.9279	146.6700	61.0603	683.6583
50.0	451.6994	144.3891	59.1048	655.1932
60.0	311.0063	130.8658	47.7834	489.6554
70.0	167.9332	108.1927	34.3338	310.4597
75.0	101.7433	88.4834	26.0710	216.2978
80.0	45.5637	58.5735	16.1263	120.2635
85.0	8.8855	12.3191	3.3810	24.5857

Table A-3. TABULATED DATA FROM SEVERAL MODELS FOR THE  
 DAVE MODEL 3 ATMOSPHERE

IO = 1353.0000  
 UW = 2.9300  
 UO = 3100  
 PR = 1013.0000  
 TAUS = .0999  
 TAU3B = .0979  
 RS = .0200  
 TEMP = 294.0000  
 CONST = .0933  
 BA = .8600  
 TAUB = .0620

ZENITH	AIRMASS
0.0000	.9995
20.0000	1.0634
30.0000	1.1536
48.1900	1.4972
50.0000	1.5525
60.0000	1.9927
70.0000	2.8997
75.0000	3.8076
80.0000	5.5790
85.0000	10.3163

## ATWATER MODEL

ZEN	TA	AW	TMD	DIRH	DIFSH	DIFGH	DTOT
0.0	.9399	.1063	.9381	983.9331	73.8571	1.4512	1059.2413
20.0	.9362	.1083	.9355	912.4993	72.0609	1.3507	985.9109
30.0	.9310	.1110	.9319	825.7991	69.6973	1.2285	896.7250
48.2	.9113	.1200	.9195	594.9837	62.1416	.9015	658.0268
50.0	.9081	.1214	.9176	567.8031	61.0969	.8628	629.7628
60.0	.8836	.1308	.9038	408.0431	54.0058	.6339	462.6827
70.0	.8348	.1465	.8795	239.3498	43.7662	.3884	283.5044
75.0	.7881	.1592	.8585	156.2876	36.7179	.2648	193.2703
80.0	.7031	.1790	.8233	78.9828	27.4474	.1460	106.5762
85.0	.5087	.2177	.7472	17.4954	14.2687	.0436	31.8077

## WATT MODEL

ZEN	TA	TH20A	TH20S	T03	TAIR	DIRH	DIFSH	DIFGH	DTOT
0.0	.9762	.9146	.9379	.9768	.9016	997.790	158.878	1.029	1157.697
20.0	.9743	.9137	.9341	.9763	.8974	926.307	154.267	.974	1081.548
30.0	.9716	.9125	.9287	.9757	.8917	839.372	148.600	.907	988.879
48.2	.9609	.9088	.9083	.9733	.8714	606.839	133.046	.725	740.610
50.0	.9592	.9083	.9051	.9729	.8683	579.335	131.160	.703	711.198
60.0	.9452	.9047	.8797	.9698	.8451	417.053	119.769	.572	537.395
70.0	.9157	.8992	.8291	.9635	.8029	244.378	106.911	.428	351.717
75.0	.8921	.8953	.7813	.9586	.7702	161.311	100.017	.351	261.680
80.0	.8457	.8896	.6933	.9489	.7142	83.046	92.861	.273	176.179
85.0	.7602	.8800	.4896	.9309	.6123	22.017	85.480	.193	107.689

## HOYT MODEL

ZEN	TAS	AA	AW	ACO2	A03	A02	TR
0.0	.9321	.0466	.1272	.0075	.0258	.0075	.9170
20.0	.9279	.0464	.1298	.0076	.0264	.0079	.9125
30.0	.9220	.0461	.1333	.0078	.0274	.0085	.9063
48.2	.9000	.0450	.1451	.0084	.0305	.0107	.8840
50.0	.8965	.0448	.1469	.0085	.0310	.0110	.8807
60.0	.8692	.0435	.1592	.0091	.0344	.0137	.8560
70.0	.8155	.0408	.1796	.0101	.0401	.0190	.8138
75.0	.7650	.0383	.1960	.0109	.0448	.0242	.7781

ZEN	DIRH	DIFSH	DIFGH	DTOT
0.0	908.3464	98.2025	1.2750	1007.8240
20.0	841.6283	97.2154	1.1857	940.0295
30.0	760.6718	95.8785	1.0773	857.6276
48.2	545.5922	91.1694	.7888	637.5504
50.0	520.3379	90.4580	.7548	611.5508
60.0	372.5084	85.1641	.5554	458.2279
70.0	218.1138	76.0972	.3449	294.5559
75.0	142.9609	68.9767	.2407	212.1782

Table A-3. TABULATED DATA FROM SEVERAL MODELS FOR THE DAVE MODEL 3 ATMOSPHERE (concluded)

LACIS MODEL

ZEN	AW	A03	DTOT
0.0	.1327	.0224	1093.1443
20.0	.1350	.0231	1020.9793
30.0	.1379	.0239	933.1311
48.2	.1475	.0270	697.4798
50.0	.1489	.0275	669.5094
60.0	.1585	.0312	503.7295
70.0	.1736	.0382	324.5709
75.0	.1849	.0448	233.1836
80.0	.2012	.0568	143.4038
85.0	.2282	.0851	59.8739

DAVIES MODEL

ZEN	DIRH	DIFSH	DIFGH	DTOT
0.0	936.3331	129.0411	1.7442	1067.1183
20.0	866.5097	127.0680	1.6427	995.2204
30.0	781.7525	124.4193	1.5186	907.6905
48.2	556.3066	115.3767	1.1823	672.8657
50.0	529.8009	114.0456	1.1419	644.9885
60.0	374.4382	104.3899	.8989	479.7270
70.0	212.2479	88.2198	.6243	301.0919
75.0	134.2798	75.3692	.4742	210.1232
80.0	64.7464	56.4768	.3127	121.5359
85.0	14.9323	28.3066	.1388	43.3777

BIRD MODEL

ZEN	TA	T03	TU	TR	TAS	AW	TAA
0.0	.9219	.9834	.9874	.9137	.9286	.1219	.9927
20.0	.9175	.9826	.9872	.9094	.9247	.1235	.9923
30.0	.9115	.9816	.9869	.9033	.9192	.1257	.9916
48.2	.8891	.9778	.9860	.8816	.8989	.1330	.9891
50.0	.8856	.9772	.9859	.8783	.8961	.1340	.9883
60.0	.8585	.9727	.9849	.8531	.8722	.1411	.9843
70.0	.8068	.9643	.9834	.8074	.8279	.1520	.9745
75.0	.7595	.9566	.9822	.7684	.7926	.1601	.9582
80.0	.6774	.9430	.9803	.7078	.7444	.1717	.9099
85.0	.5057	.9116	.9770	.6157	.7804	.1907	.6479

ZEN	DIRH	DIFSH	DIFGH	DTOT
0.0	938.8409	74.5702	1.6248	1035.0359
20.0	871.3803	93.1824	1.5273	966.0901
30.0	789.4391	91.3142	1.4082	882.1616
48.2	571.0358	85.0776	1.0865	657.1999
50.0	545.2946	83.9642	1.0470	630.3058
60.0	393.9366	77.2467	.8155	471.9989
70.0	234.2099	65.8789	.5567	300.6456
75.0	155.8315	55.7656	.4135	212.0107
80.0	83.3321	39.5922	.2569	123.1812
85.0	25.5650	11.1038	.0729	36.7417

<b>Document Control Page</b>	1. SERI Report No. TR-642-761	2. NTIS Accession No.	3. Recipient's Accession No.
4. Title and Subtitle  A Simplified Clear Sky Model for Direct and Diffuse Insolation on Horizontal Surfaces	5. Publication Date February 1981		6.
	7. Author(s) Richard E. Bird, Roland L. Hulstrom		8. Performing Organization Rept. No.
9. Performing Organization Name and Address Solar Energy Research Institute 1617 Cole Boulevard Golden, Colorado 80401	10. Project/Task/Work Unit No. 1093.00		11. Contract (C) or Grant (G) No. (C) (G)
	12. Sponsoring Organization Name and Address		13. Type of Report & Period Covered Technical Report 14.
15. Supplementary Notes			
16. Abstract (Limit: 200 words) A detailed comparison was made between five simple broadband models for clear sky global insolation. Compared models were those of Atwater and Ball, Davies and Hay, Watt, Hoyt, and Lacis and Hansen. A sixth simple model, called the Bird model, has been formulated by using parts of these five models and by comparison with the results from three rigorous radiative transfer codes. All of the simple models provide results that agree within <10% with the three rigorous codes when the sun is in the zenith position. The Bird and Hoyt models agree within 3% with each other and with the results of the rigorous codes. However, the Bird model is easier to implement and has broader application than the Hoyt model.			
17. Document Analysis a. Descriptors Comparative Analysis ; Data ; Energy Models ; Insolation:T1 ; Mathematical Models:Q1 ; Measuring Methods ; Meteorology ; Solar Radiation ; Radiations  b. Identifiers/Open-Ended Terms    c. UC Categories 59,61,62,63			
18. Availability Statement National Technical Information Service U.S. Department of Commerce 5285 Port Royal Road Springfield, Virginia 22161		19. No. of Pages 53 20. Price \$5.25	